

To appear in The Astrophysical Journal

## Spatially Resolved Far-Ultraviolet Spectroscopy of the Nuclear Region of NGC 1068 <sup>1</sup>

Wei Zheng<sup>2</sup>, Jun-Xian Wang<sup>2,3</sup>, Gerard A. Kriss<sup>2,4</sup>, David Sahnou<sup>2</sup>, Mark Allen<sup>5</sup>, Michael Dopita<sup>6</sup>, Zlatan Tsvetanov<sup>2,7</sup>, and Geoffrey Bicknell<sup>6</sup>

### ABSTRACT

We carry out high-resolution FUSE spectroscopy of the nuclear region of NGC 1068. The first set of spectra was obtained with a 30'' square aperture that collects all emission from the narrow-line region. The data reveal a strong broad O VI component of FWHM  $\sim 3500 \text{ kms}^{-1}$  and two narrow O VI  $\lambda\lambda 1031/1037$  components of  $\sim 350 \text{ kms}^{-1}$ . The C III  $\lambda 977$  and N III  $\lambda 991$  emission lines in this spectrum can be fitted with a narrow component of FWHM  $\sim 1000 \text{ kms}^{-1}$  and a broad one of  $\sim 2500 \text{ kms}^{-1}$ . Another set of seven spatially resolved spectra were made using a long slit of  $1''.25 \times 20''$ , at steps of  $\sim 1''$  along the axis of the emission-line cone. We find that (1) Major emission lines in the FUSE wavelength range consist of a broad and a narrow component; (2) There is a gradient in the velocity field for the narrow O VI component of  $\sim 200 \text{ kms}^{-1}$  from  $\sim 2''$  southwest of the nucleus to  $\sim 4''$  northeast. A similar pattern is also observed with the broad O VI component, with a gradient of  $\sim 3000 \text{ kms}^{-1}$ . These are consistent with

---

<sup>1</sup> Based on observations made with the NASA-CNES-CSA Far Ultraviolet Spectroscopic Explorer (FUSE), which is operated for NASA by the Johns Hopkins University under NASA contract NAS5-32985, and observations with the NASA/ESA *Hubble Space Telescope*, obtained at the Space Telescope Science Institute, which is operated by the Association of Universities of Research in Astronomy, Inc., under NASA contract NAS5-26555.

<sup>2</sup>Center for Astrophysical Sciences, Department of Physics and Astronomy, The Johns Hopkins University, Baltimore, MD 21218

<sup>3</sup>Center for Astrophysics, University of Science and Technology of China, Hefei, Anhui, 230026, China

<sup>4</sup>Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218

<sup>5</sup> Observatoire de Strasbourg, 67000 Strasbourg, France

<sup>6</sup> Mount Stromlo & Siding Spring Observatories, Australian National University, ACT 2611, Australia

<sup>7</sup>NASA Headquarters, Washington, DC 20546-0001

the HST/STIS findings and suggest a biconical structure in which the velocity field is mainly radial outflow; (3) A major portion of the C III and N III line flux is produced in the compact core. They are therefore not effective temperature diagnostics for the conical region; and (4) The best-fitted UV continuum suggests virtually no reddening, and the He II 1085/1640 ratio suggests a consistently low extinction factor across the cone. At  $\sim 2''$  northeast of the nucleus there is a region characterized by (a) a strong Ly $\alpha$  flux, but normal C IV flux; (b) a broad O VI line; and (c) a significantly enhanced C III flux.

*Subject headings:* galaxies: individual (NGC 1068) — galaxies: Seyfert galaxies: active —

## 1. INTRODUCTION

NGC 1068 is a prototypical Seyfert 2 galaxy. Because of its proximity ( $z = 0.0038$ ) and brightness, it has been studied in nearly every possible detail. The polarimetric observation by Antonucci et al. (1985), which reveals a Seyfert 1 spectrum in scattered light, suggests that the nucleus and its associated broad-line region (BLR) are obscured. This finding provides strong evidence for the unified theory in which viewing angles account for the differences between various active galactic nuclei (Antonucci 1993)

The nuclear region of NGC 1068 harbors a variety of astrophysical phenomena. At the very center of the nucleus there is a bright compact ( $< 0''.3$ ) region commonly referred as “the hot spot”. Within a few arcseconds from the nucleus, there are several bright and compact clouds that coincide with knots in the radio jets (Wilson & Ulvestad 1983; Evans et al. 1991). The narrow-line region (NLR) is conical in shape toward the northeast (Fig. 1), along a position angle of  $\sim 200$  deg and with an opening angle of  $\sim 40$  deg. Beyond a  $6''$  radius the surface brightness drops dramatically, and emission is dominated by two ring-like filaments at  $\sim 10''$  and  $15''$  from the nucleus.

High-spatial-resolution spectroscopy of NGC 1068 has been carried out in the optical (Caganoff et al. 1991; Unger et al. 1992; Inglis et al. 1995; Emsellem et al. 2005; Gerssen et al. 2006) as well as in the UV (Caganoff et al. 1991; Kraemer, et al. 2000a; Crenshaw & Kraemer 2000b; Cecil et al. 2002; Groves et al. 2004). From approximately  $2''$  southwest of the nucleus to  $4''$  northeast, emission lines exhibit multiple components (Cecil et al. 1990; Crenshaw & Kraemer 2000a): (1) major emission lines consist of narrow and broad lines; (2) broad lines are approximately  $2500\text{--}4000 \text{ km s}^{-1}$  wide, which may be linked to those that are found in polarized light and believed to be reflected light from the inner BLR, and (3) narrow lines consist of a pair

of red and blue components. The [O III] and [N II] line profiles suggest that the separation of these two components varies across the conical NLR. In addition to an overall biconical ionization configuration, there are compact knots whose optical spectra resemble kinematically the associated absorption line systems in quasars (Cecil et al. 2002; Crenshaw, et al. 2003; Crenshaw & Kraemer 2005). These line-emitting knots have blueshifted radial velocities up to  $3000 \text{ km s}^{-1}$  relative to the galaxy’s systemic velocity, contributing mostly to the emission-line flux but not the continuum. Between  $\sim 2''.5$  and  $4''.5$  northeast from the nucleus, UV line emission is redshifted relative to the systemic value, a pattern that is interpreted as the expansion of the plasma in the radio lobe (Axon et al. 1998).

Several important emission lines in the far-UV (FUV) region between 912 and  $1150 \text{ \AA}$  are observable only with specially crafted UV instruments. During the Astro-1 mission the Hopkins Ultraviolet Telescope (HUT) observed NGC 1068 with  $18''$  and  $30''$  apertures. The most striking features in the wavelengths below  $1150 \text{ \AA}$  are the strong C III  $\lambda 977$  and N III  $\lambda 991$  lines. The line intensity ratios of C III I( $\lambda 909$ )/I( $\lambda 977$ ) and N III I( $\lambda 1750$ )/I( $\lambda 991$ ) are temperature sensitive, and the derived temperature is  $> 25\,000 \text{ K}$  (Kriss et al. 1992), higher than the values expected for a region producing C III and N III emission by photoionization. The line ratios in NGC 1068 are similar to those of the Cygnus Loop supernova remnant (Blair et al. 1991), suggesting a significant contribution from shock-heating mechanisms. Astro-2 observations of NGC 1068 were obtained with a  $12''$  aperture at three different positions. The results (Grimes et al. 1999) suggest that the emission lines observed with HUT likely arise in the inner nuclear region imaged with HST. However, the poor angular resolution of the HUT instrument ( $> 10''$ ) does not allow a study of the NLR in terms of spatial details.

The line-emitting mechanisms in the NLR, *i.e.* photoionization from the nucleus or shocks produced by jets, have long been under debate. Dopita & Sutherland (1996) and Bicknell et al. (1998) proposed that emission in the NLR may be entirely caused by shocks. Velocity splitting over  $1000 \text{ km s}^{-1}$ , reported by Axon et al. (1998), in the vicinity of some of the bright emission-line knots provides evidence that fast shocks exist in the NLR of NGC 1068. However, Seyfert galaxies host powerful nuclear sources of ionizing radiation, and the situation can be more complex with shocks or nuclear photoionization prevailing in different environments (Allen et al. 1998; Morganti et al. 1998). Morse et al. (1996) suggested that photoionizing shocks are important when a radio jet interacts with the interstellar medium, but not in the objects where sharp, straight-edged ionization cones are observed. Ferguson et al. (1995) argued that strong C III  $\lambda 977$  and N III  $\lambda 991$  emission may arise from fluorescence in photoionized gas if turbulent velocities exceed  $\sim 1000 \text{ km s}^{-1}$ . However, Grimes et al. (1999) found that such a velocity would lead to extreme physical conditions that are inconsistent with the Astro-2 data. More recent HST data (Crenshaw & Kraemer

2000b; Cecil et al. 2002) found that the emission-line ratios are consistent with photoionization instead of shock heating mechanism.

We have carried out observations with the *Far Ultraviolet Spectroscopic Explorer* (FUSE) to study the spatial distribution of FUV emission lines. In this paper we present the results of both large-aperture and spatially resolved spectroscopic observations with FUSE. For the first time, we are able to study the position dependence of several important diagnostic lines in the FUV band.

## 2. DATA

### 2.1. FUSE data

FUSE covers a wavelength range between 904 and 1188 Å, with a spectral resolution of  $R \sim 20000$  (Moos et al. 2000). It is based on a Rowland circle design and consists of four separate optical paths or channels. Each channel consists of a mirror coated with aluminum plus LiF or SiC, a focal plane assembly, which includes the spectrograph apertures, a diffraction grating coated with aluminum plus LiF or SiC, and a portion of FUV detector (two named A and B). In all, data are collected from eight traces, and the pairs of A and B are called channels. Our original plan was to take spectra of the nuclear region of NGC 1068 at seven positions separated by  $\sim 1''$ . Because of thermal instability of the optical structure in orbit, the four channels in the FUSE spectrograph are not perfectly aligned, with orbit-dependent drifts possibly as large as  $6''$ . Therefore, a new observing strategy was developed to counter the drifting effect. The high-resolution aperture (HIRS:  $1''.25 \times 20''$ ) is used for spectroscopy on the finest spatial scales, to ensure that maximum resolution is maintainable even if the telescope imaging or pointing stability degrades below specifications. Since the LiF1 channel is mounted on the same optical system as the FUSE fine error sensor, the spatial knowledge and stability is fully available for the data from this channel.

The observations of NGC 1068 were carried out with two modes between 2001 November 28 and December 02. The first set of data of 21951 seconds was taken with fixed pointing and the LWRS (low-resolution) square aperture of  $30''$ . With such a large aperture, fluxes of the nuclear region are all collected (Fig. 2), even with the thermally-induced mirror motions anticipated. The second set of observations was carried out in a scanning mode across the conical emission-line region, with a total exposure of 94434 seconds. A narrow HIRS slit at a position angle of 137 deg was used. Because of orbital constraints, we were not able to place the slit completely perpendicular to the conical axis. The slit actually makes an angle of  $\sim 97$  deg with respect to the conical axis.

Over years of FUSE operation, a large amount of engineering data have been collected, and the drift pattern among the four channels has been well studied. After peak-up alignments of channels with a bright point source at the beginning of orbital nights, the relative drifts exhibit a dependence on orbital time (Fig. 3). The LiF2 channel exhibits relatively small drifts, within one arcsecond with respect to the LiF1 channel. The SiC channels exhibit drifts that can be represented by a constant drift rate during orbital nights. For example, the SiC2 channel drifts relative to the LiF1 channel at a rate of approximately  $4''$  per 30 minutes. The starting drift at the orbital sunset is however variable, depending on the degree of occultation because the drift is generally in the reverse direction during orbital days.

NGC 1068 itself is too diffuse to appear in the guide-star camera. For target acquisition, we used an offset guide star  $\sim 85''$  from the nucleus. We also used a FUV-bright star, Feige 23, which is approximately four degrees away from the source, for instrumental peak-up alignments at the beginning of each pointing. One orbit of peak-up operation on Feige 23 was made to establish the drift pattern due to orbital motion. During the second orbit, one peak-up with the MDRS aperture ( $4'' \times 20''$ ) and another with HIRS were carried out on Feige 23. The instrument was then pointed to the southwest part of the nucleus of NGC 1068, and we started observations in a scanning mode. As shown in Table 1, each of the five pointings consists of 7-9 orbits on the target. The observation window varied orbit by orbit, between  $\sim 1200$  and  $2600$  sec. The scanning speed in each individual orbit was set to complete  $10''$  within the designated observation window. After accumulating a total exposure time of  $\sim 18000$  sec, FUSE returned to Feige 23 and carried out one peak-up with MDRS and another with HIRS, then moved back to NGC 1068 for a new set of scans. After a total of five pointings, FUSE returned to Feige 23 for one last orbit of peak-up with MDRS, to acquire additional information on the drift pattern.

Since the channel alignments are not perfect except at the beginning of the first orbit, the orbit-time dependence of channel drifts derived from the engineering data is only useful in terms of the relative drift speed, not for the absolute timing.

The drifts in the SiC channels can be as large as  $6''$ , which is near the size of the ionization cone. The drift direction is mainly along  $+x$  on the FUSE aperture plate, i.e., along the conical direction of NGC 1068 at the desired FUSE aperture position angle of  $\sim 105^\circ$ . Therefore the observation procedure was designed so that each observation starts  $\sim 5''$  from the nucleus in the anti-cone (southwest) direction and sweeps across the cone towards the northeast direction at a constant angular velocity. Fig. 4 illustrates the time sequence in one orbit of observation: with a constant angular speed, the FUSE HIRS slit swept across the nuclear region of NGC 1068. We identified the peak position of the count rate in the LiF1 channel as the nuclear position. Based on the angular speed illustrated in Fig. 3, seven time

intervals were defined, and the spectral extraction in these segments yields the data A-G. The upper panel of Fig. 4 marks the segments for the LiF1 channel, whose spatial information is fully available. For the other three channels, their count rates peaked at different times as a result of drifting. We marked their respective peak positions. The sweeping speeds in these channels are the sum of two terms: that of the LiF1 channel plus the relative drift speed of each channel (Fig. 3) with respect to LiF1. Their time segments A-G are also illustrated in Fig. 4. The observation results in a series of segments that are dependent on the relative positions with respect to the nucleus. In each time sequence, one can determine the time when the observation slits coincide with the nucleus, as determined by the peak in the count rate.

The LiF1 slit positions are illustrated in Fig. 1. Seven sets of data are extracted with a relative shift of 1 arcsec ( $1''.25$  at the nucleus), starting from  $2''$  southwest from the nucleus (in the opposite direction of the cone), to  $\sim 4''$  northeast of it. They were named as A,B,C (the nucleus, see Fig. 1), D,E,F and G. The significant astigmatism of FUSE yields a spatial resolution of  $\sim 5''$  at best, therefore it cannot be used like a normal long-slit spectrograph to dissect the fine details along its slit. Not all the segment data are extracted: If a time segment is incomplete (“G” in Fig. 4, for example), they are not used.

Four pairs of spectra were derived from the FUSE channels. Each pair of “A” and “B” data is subject to the same normalization factor. We used the O VI emission flux, which is common to all four channels, to normalize the spectra to the level in LiF1A. We first calculated the continuum level between 1050 and 1070 Å for LiF2B, SiC1A, SiC 2B and LiF1A, then subtracted it from the spectrum to enable the measurement of emission-line fluxes. We then calculated the O VI flux in two bins: a narrow component between 1041.1 and 1043.6 Å, and a broad component between 1027 and 1048 Å. The flux ratios with respect to their values in the LiF1A spectrum were calculated, then their mean values were used as the normalization factors (Table 2). For the first set of data, taken with LWRS, we corrected the “worm effect” in the LiF1A,B channel, according to the procedures described in the *FUSE Instrument and Data Handbook* (<http://fuse.pha.jhu.edu/analysis/dhbook.html>).

FUSE data reduction was carried out with pipeline version CALFUSE2.3, with specific timing flags to extract time-dependent (hence position-dependent) segments of spectra in eight traces. The FUSE spatial resolution is  $1''.5$ , and the spacecraft jitter, as determined from the FUSE engineering data, is approximately  $0''.6$ . The total spatial resolution is therefore  $1''.6$ . Since the data were obtained with a  $1''.25$  slit, there is overlap between positions. This can be seen by the fact that the sum of the fluxes in all positions is larger than that taken with one large  $30''$  aperture.

The normalized spectra in the four channels are binned to  $0.1\text{Å}$ , then merged, weighted

by their signal-to-noise ratio. The merged spectra at the seven slit positions are plotted in Fig. 5. The wavelength bin between  $\sim 1077$  and  $1087 \text{ \AA}$  falls into a gap between the LiF channels, and only data in the SiC2B channel exist.

## 2.2. HST data

We retrieved archival HST/STIS spectroscopic data to complement our FUSE data. Two sets of UV spectra are available: one taken at a position angle of  $218^\circ$  and four adjacent slit positions of  $0''.2 \times 52''$  (Cecil et al. 2002, STIS-A hereafter), and another at a position angle of  $202^\circ$  taken with a slit of  $0''.1 \times 52''$  (Crenshaw & Kraemer 2000b, STIS-B). Both sets (see Fig. 1) contain data taken with grating G140L and G230L, covering a wavelength range of  $\sim 1150$  to  $3170 \text{ \AA}$  at a resolution of  $R \sim 1000$ . These STIS observations were made between 1988 and 2000, therefore not simultaneous with the FUSE data. For STIS-A, we only used three slit positions as the other does not have G140L data. The fluxes in the three extracted spectra are summed up.

Since the FUSE spatial resolution ( $\geq 5''$  along its slit) is considerably lower than that of STIS, and since the position angles of these two sets of observations are nearly perpendicular, we must normalize the STIS data to that of FUSE. Using the continuum level around  $1180 \text{ \AA}$  in these spectra did not result in satisfactory matches as the line/continuum ratios are not constant at these spatial scales across the nucleus. We therefore use HST WFPC2 images retrieved from MAST to determine the normalization factor between STIS and FUSE data.

To extract the STIS spectra, we first smoothed the 2-D images by 30 pixels ( $0''.67$ ) with a Gaussian kernel along the spatial direction of the FUSE scans, to match the spatial resolution of the corresponding FUSE data. We then ran the standard STIS pipeline task “x1d” with steps of one arcsecond ( $1''.25$  at the nucleus) that correspond to the respective moving FUSE slit positions, and a slit width of  $1''.25$ . To normalize the STIS fluxes, we used a HST/WFPC2 image of NGC 1068, taken with filter F218W, and smoothed it along the same direction with the same kernel size as the STIS 2-D image. We then measured the flux in each STIS window. To normalize the FUSE fluxes, we smoothed the WFPC2 image to the FUSE resolution, namely  $0''.67$  and  $5''$ , respectively, along the direction perpendicular and parallel to that of the FUSE slit (Fig. 1). The flux ratios in respective extraction windows are used to derive the normalization factors (Table 2). To the northwest direction from the source, there are several bright spots whose fluxes may be picked up because of FUSE’s low spatial resolution along its slit. We tested with several extraction windows that are sufficiently short not to include the fluxes from these bright spots in the smoothed WFPC2 image. Since it is unlikely that these bright spots contribute to the redshifted UV emission

lines, we used the flux values measured in a short window in calculating the scaling factor. Any uncertainties introduced by this window selection only affect the scaling of Ly $\alpha$  and C IV in position A and G.

Since the FUSE slit positions are accurate to  $\sim 0''.3$ , we measured the fluxes in the neighboring regions, to make sure that the normalization factors are accurate to at least 20%. The height of STIS extraction windows is the same as that of FUSE ( $\sim 1''$ ), but their widths ( $0''.1$ - $0''.2$ ) are much smaller than FUSE ( $20''$ ). Therefore considerable STIS normalization factors were applied. The matching STIS spectra are plotted in Fig. 6.

We compared extracted spectra from the STIS-A and STIS-B datasets, and their line ratios are comparable within 25%. In the following section, we only use the data from STIS-A, as they cover a spatial region six times as wide ( $0''.2 \times 3''$ ) as that of STIS-B.

### 3. FITTING

Spectral analyses were carried out using IRAF task *Specfit* (Kriss 1994). For the FUSE spectrum taken with a large aperture, we used a pair of narrow components and one broad component in the O VI emission line profile. Each narrow O VI component was modeled as a doublet whose wavelength ratio is fixed by atomic data and whose line widths are identical. The O VI emission feature is heavily absorbed by interstellar absorption at  $\sim 1037\text{\AA}$ , therefore we introduced several absorption components. The Ly $\beta$  emission is not prominent, and it is often overwhelmed by the broad O VI emission. Only in one or two positions is the Ly $\beta$  emission visible. We therefore only modeled one component with a fixed line width. For N III and C III emission line features, one narrow component and one broad component were included in fitting. We fitted a power law with no extinction to the continuum. The fitted results are listed in Table 3.

For the FUSE spectra taken at positions A-G, we used one narrow component plus one broad component to model the O VI emission, and one component for the N III and C III lines. To reduce the effect of channel drifts, the FUSE spectra were fitted individually for each channel. The fitting results in Table 4 are primarily derived from the LiF1 data, which are not affected by channel drifts. For wavelengths below  $995\text{\AA}$ , we used the fitting results from the scaled SiC2A channel, which extends to  $1005\text{\AA}$  and is therefore less susceptible than the SiC1B channel to the detector edge effects. For He II 1085, LiF2A data were used. The fluxes of fitted components were normalized by the values in Table 2, and then tabulated in Table 4. The profiles of four major emission lines at seven different slit positions are plotted in Fig. 7, to show the changes in the narrow and broad components as a function of spatial



location.

The UV spectra suggest a low extinction level. We first fitted the continuum in the wavelength windows that are free of emission and absorption lines, and the results suggest a power-law index of  $\alpha = 0.90 \pm 0.04$  ( $f_\lambda \propto \lambda^\alpha$ ) and an extinction of  $E_{B-V} = 0.005 \pm 0.002$ . As a comparison, the HUT spectrum between 912 and 1800 Å was fitted with a power-law continuum of  $\alpha = 0.62 \pm 0.15$  and  $E_{B-V} = 0.065 \pm 0.02$ . If we adopt this extinction value, the power-law index is  $\alpha = 0.85 \pm 0.06$ . The similar power indices suggest that one single power-law continuum can fit the far-UV spectrum of NGC 1068 without introducing significant reddening, as suggested from the UV/optical line ratios (Crenshaw & Kraemer 2000a).

## 4. Discussion

Many previous studies have gradually unraveled the kinematical complexities of gas in the nuclear region of NGC 1068. The X-ray data (Kinkhabwala et al. 2002) suggest that emission lines are formed mainly in a photoionized plasma of a temperature around a few eV. The spatial resolution afforded by HST leads to a picture of decelerating jet (Das et al. 2006): a biconical outflow from the nuclear region sweeps up denser, ambient clouds in the interstellar medium (ISM) of NGC 1068. Other possibilities include overlapping, discrete ejection that gradually dissipates (Axon et al. 1998; Capetti et al. 1997). In addition, a high-velocity radio jet impinges on some of the clouds. Some gas expands perpendicularly to the axis of the jet, and the expanding radio lobe at the end of the jet also pushes on the ambient ISM. Near the nucleus, kinematic components span several thousand kilometers per second in velocity, and the continuum hot spot visible in HST images reflects a polarized view of the broad lines in the active nucleus.

Our spatially resolved FUSE observations, while not at the resolution of HST, add information from major emission lines shortward of the HST bandpass at high spectral resolution. We plot the fluxes of major UV emission lines from our observations at seven slit positions in Fig. 8, and list the fitted line properties in Table 4. In this section, we discuss the four emission lines in the FUSE spectral range, along with a comparison to emission lines observed in the STIS spectra.

### 4.1. O VI emission

The most prominent feature in the FUSE spectra is the O VI  $\lambda\lambda 1031, 1037$  emission line. In the data taken with LWRS, the two narrow O VI components are of FWHM  $\sim 350$

$\text{km s}^{-1}$  and a separation of  $\sim 200 \text{ km s}^{-1}$ . In the seven spectra taken with a narrow slit, we resolve this blend into one narrow component with FWHM of  $\sim 350 \text{ km s}^{-1}$  and one broad component. In Fig. 9 we plot the O VI profiles at the seven different slit positions. The high spectral resolution of FUSE data enables us to compare with the results of optical Fabry-Perot spectroscopy (Cecil et al. 1990), which reveal a narrow core of  $\sim 300 \text{ km s}^{-1}$ . According to the optical data, nearly 75% of the [N II]  $\lambda 6583$  flux is from components of  $\sim 1500 \text{ km s}^{-1}$  wide. Line widths at such a scale are consistent with that derived from HST UV spectroscopy. The narrow-line flux in the FUSE spectra is highly concentrated (60%) at the compact core, suggesting that the “true” NLR probably remains unresolved, at a sub-arcsecond scale. While the narrow O VI emission line is not the dominant component, its distribution is different from its broad counterpart, while only 40% of the broad O VI line flux is from slit position C.

It is surprising that the O VI emission is dominated by a component that is broader than those seen in polarized light: in the LWRS spectrum, more than 3/4 of the O VI flux is from a component with FWHM  $\sim 3500 \text{ km s}^{-1}$  that is blue-shifted relative to the narrow component by  $\sim 500 \text{ km s}^{-1}$ . This broad component is present in all seven FUSE slit spectra with considerable strength. It may arise from the reflected emission from the hidden BLR, and/or may be the result of a significant velocity dispersion in the NLR. The HST FOC data (Axon et al. 1998) reveal that emission lines near the hot knots 2" northeast (FUSE slit positions D and E) are split into two velocity systems separated by  $\sim 1500 \text{ km s}^{-1}$ . The STIS spectra discussed by Groves et al. (2004) show [O III] emission knots spanning such a broad velocity range in the immediate vicinity of the nucleus, but not at distances of several arcseconds. Since the FUSE slit collects emission from a block of regions spanning several arcseconds perpendicular to the conical axis, the total line emission from these regions may be blended into one broad component.

The ratios of C IV/O VI may provide insight into the physical conditions of the line-emitting regions. The value is higher for the narrow components than their broad counterparts, implying a range of the ionization parameter  $U \sim 0.05 - 1.0$  in a typical photionization calculation. High values of  $U > 1$  are consistent with models that assume the same origin for the associated absorbers and BLR (Kriss et al. 2003), suggesting that the clouds that produce the broad emission components may be of the same origin as associated absorbers.

## 4.2. Velocity field

The narrow O VI line exhibits a systematic velocity shift from position A to G by approximately  $220 \text{ km s}^{-1}$ . This gradient in the spatially resolved spectra explains why

there are two narrow-line components in the integrated flux from the large-aperture FUSE spectrum, where it is unresolved. Crenshaw & Kraemer (2000a) reported a similar velocity pattern in their HST spectra. Das et al. (2006) successfully modeled this as a biconical outflow in which radial velocity changes as a function of the distance to the central nucleus: the emission line knots show evidence for radial acceleration to a projected distance of  $2''$  to the northeast direction, followed by deceleration up to  $4''$ . The O VI line widths also increase at  $\pm 2''$  from the nucleus, probably implying a larger dispersion in these regions.

The broad component of O VI exhibits a qualitatively similar kinematic pattern, but at a larger amplitude: its centroid shifts by  $\sim 1500 \text{ km s}^{-1}$  (Fig. 10) across the same spatial region. Fig. 11 shows similar trends for the narrow and broad components of C IV in the HST spectra which have not been explicitly noted in previous studies: the narrow component follows the kinematic pattern of the optical lines modeled by Das et al. (2006); the broad component of C IV shows behavior similar to that of O VI in the FUSE spectra. Prior observations that noted this blue-shifted broad component invariably attributed it to the reflection of the BLR. The blue shift and width (at the position of the hot spot, and in integrated light) are comparable to the broad polarized  $H\beta$  line observed by Antonucci et al. (1985).

In a scattered BLR picture, a blue shift is caused by the outflowing wind from the torus along our line of sight, and a line width is due to the intrinsic broad line width convolved with the thermal width of the hot reflecting wind. As a broad component is present in the FUSE spectra at all seven positions, it is possible to assume that this is reflected light from the hidden BLR. However, a large covering factor is needed to explain the observed fluxes. Assuming a covering factor of 0.1, the intrinsic flux of the broad O VI emission in NGC 1068 would exceed that in NGC 4151. A more reasonable explanation for the observed broad line widths is the large velocity dispersion between bright knots. At approximately  $2''$  from the core (position A and E), the FWHMs of broad components are the broadest at  $3200 \text{ km s}^{-1}$ . These maxima coincide with the widest splitting of velocity in bright knots Crenshaw & Kraemer (2000a). With an intrinsic dispersion of  $\sim 800 \text{ km s}^{-1}$  and a separation in velocity of  $\sim 2500 \text{ km s}^{-1}$  between bright knots (from STIS results), data collected by the long FUSE slit would exhibit a broad component of  $\sim 3200 \text{ km s}^{-1}$ , which is what observed at positions A and E. Extended regions of hot, photoionized gas are seen in X-ray images of NGC 1068 (Young, Wilson & Shopbell 2001; Kinkhabwala et al. 2002) that could be visible manifestations of this hot outflow. It is natural to assume that these high-velocity clouds may be related to the associated absorbers (Crenshaw, et al. 2003, and references therein) in AGN, which are mostly blueshifted. As with the lower-velocity, lower ionization emission-line gas, the acceleration of this high ionization gas eventually is brought to a halt by an unknown deceleration mechanism, which might plausibly be interaction with the ambient ISM of NGC

1068. The evidence for deceleration at arcsecond scale may suggest that acceleration of the outflow materials may take place at sub-arcsecond scales.

### 4.3. C III $\lambda 977$

The flux ratio of C III  $\lambda 1909$  to 977 is extremely sensitive to temperature, and the value measured in N1068 suggests a high temperature that is consistent with shock-heating (Kriss et al. 1992). The FUSE data taken with a large aperture (Table 3) reveal that this emission line consists of a narrow and a broad component. The broad component is weak and hence cannot be well separated in the data segments taken with a narrow FUSE slit. The STIS data reveal that C III  $\lambda 1909$  emission can be fitted with a narrow and a broad component of FWHM  $\sim 900$  and  $3500 \text{ km s}^{-1}$ , respectively. In principal, the ratio of C III  $\lambda 1909$  to 977 should be calculated only between the narrow components.

As shown in Fig. 8, the flux of C III  $\lambda 977$  is highly concentrated in position C and D. High temperatures implied by this line emission may therefore be associated with the compact core. In a large portion of the ionization cone, the line ratios ( $\lambda 1909/\lambda 977$ ) are considerably higher, suggesting a lower temperature. However, the line ratios are considerably lower than that derived from the HUT data, where broad C III  $\lambda 1909$  component was included in the calculation.

### 4.4. N III $\lambda 991$

The line ratio of N III  $\lambda 1750$  to  $\lambda 991$  is also temperature dependent, and its value has also been used to derive a high temperature in the ionization cone. The FUSE data also reveal a pair of components in the data taken with a large aperture. Unlike C III  $\lambda 977$ , the broad component of N III  $\lambda 991$ , as shown in Table 3, is stronger than its narrow counterpart. The FUSE data taken with a narrow slit shows that nearly a half of this broad component is in position C. In other positions, the N III  $\lambda 991$  emission can only be fitted with one component of FWHM  $\sim 1000 \text{ km s}^{-1}$ . In combination with the trend for C III  $\lambda 977$ , we conclude that a bulk of flux in these temperature-sensitive emission lines is from the compact core of NGC 1068, and their intensity is not directly tied to the physical conditions in the ionization cone.

#### 4.5. He II 1085

The distribution of He II  $\lambda 1085$  is different from other emission lines in the FUSE spectra: as shown in Fig. 8, the He II flux varies smoothly across the NLR region, like that of  $\text{Ly}\alpha$ , C IV and other lines in the STIS spectra. Since He II emission is believed to be produced mainly by recombination and is insensitive to the gas temperature, this pattern of variation may simply reflect the distribution of NLR gas across the ionization cone. The ratio of He II emission  $\lambda 1640$  to  $\lambda 1085$  is believed to be a reddening indicator. As shown in Fig. 12, the ratio is nearly constant ( $\sim 5.5$ ) between slit positions B and F. This is consistent with the HUT result of  $5.8 \pm 1.6$ , suggesting an insignificant level of extinction ( $E_{B-V} \leq 0.05$ ). The low values at positions A and G may suggest that the data at these positions are not reliable.

The He II I(1640)/I(1085) ratio is slightly lower than that expected from recombination, and this may suggest that the He II  $\lambda 1085$  may be slightly contaminated by other weak UV emission lines. N II  $\lambda 1085$  is a likely candidate, and the trend of decreasing He II (1640)/I(1085) ratio with distance from the nucleus supports this as there is a noticeable decrease in ionization state at larger distances from the nucleus (Axon et al. 1998; Kraemer, et al. 2000a; Kraemer & Crenshaw 2000b). As with O VI and C IV, there appears to be a broad component that is considerably blueshifted. However, the relevant wavelength range is in a gap between the LiF channels, and only SiC2B data are available. Because of the low S/N ratio, the reality of such a broad component is still questionable, as no such counterpart is found in the corresponding HST spectra of He II  $\lambda 1640$ .

### 5. Summary

We have carried out high-spectral resolution, spatially resolved spectroscopy of the nuclear region of the Seyfert 2 galaxy NGC 1068. The high spectral resolution of FUSE data enables us to study the line profiles of O VI, N III, and C III as a function of position across the nuclear region. Our observations using a long slit of  $1''.25 \times 20''$  at steps of  $\sim 1''$  along the axis of the emission-line cone resulted in a set of seven spatially resolved spectra running from  $\sim 2''$  southwest of the nucleus to  $\sim 4''$  northeast.

The O VI profiles exhibit considerable structure: a prominent broad component with FWHM  $\sim 3500 \text{ km s}^{-1}$ , and a narrow component of  $\sim 300 \text{ km s}^{-1}$ . Both components show a position-dependent velocity gradient along the conical axis with a velocity shift of  $\sim 220 \text{ km s}^{-1}$  in the narrow component and a shift of  $\sim 1500 \text{ km s}^{-1}$  in the broad component. Both patterns are consistent with radial outflow from the nuclear region. Both the continuum and

emission lines suggest low extinction: the UV continuum is flat to the shortest wavelengths in the FUSE spectrum, and the best fit is a power law with virtually no reddening. The He II 1640/1085 ratio also is reflective of a consistently low extinction factor across the emission-line cone. The majority of the C III and N III emission arises in the compact core, suggesting that the region with extremely high temperature is very close to the nucleus and remains unresolved. The line emission at  $\sim 2''$  northeast of the nucleus is strong and broad, characterized by (a) a strong Ly $\alpha$  flux, but normal C IV flux; (b) a broad O VI line; and (c) a significantly enhanced C III flux, possibly the result of hot knots associated with shock heating.

W.Z. would like to thank the FUSE team, particularly W. Blair, for their painstaking help in planning the observations of NGC 1068 and post-observation studies of engineering data.

This work has been supported in part by NASA grant NAG-8-1527 and NAG-8-1133. The FUSE and HST data presented in this paper were obtained from the Multimission Archive at the Space Telescope Science Institute (MAST).

## REFERENCES

- Allen, M. G., Dopita, M. A., & Tsvetanov, Z. I. 1998, *ApJ*, 493, 571
- Antonucci, R. 1993, *ARAA*, 31, 473
- Antonucci, R. R. J., & Miller, J. S. 1985, *ApJ*, 297, 621
- Axon, D. J., Marconi, A., Capetti, A., Machetto, F. D., Schreier, E., & Robinson, A. 1998, *ApJ*, 496, L75
- Bicknell, G. V., Dopita, M. A., Tsvetanov, Z. I., & Sutherland, R. S. 1998, *ApJ*, 495, 680
- Blair, W. P., et al. 1991, *ApJ*, 379, L33
- Caganoff, S., et al. 1991, *ApJ*, 377, L9
- Capetti, A., Axon, D. J., & Macchetto, F. D. 1997, *ApJ*, 487, 560
- Capetti, A., Axon, D. J., Machetto, F. D., Sparks, W. B., & Boksenberg, A. 1996, *ApJ*, 469, 554
- Cecil, G., Bland, J., & Tully, R. B. 1990, *ApJ*, 355, 70

- Cecil, G., Dopita, M. A.; Groves, B., Wilson, A. S.; Ferruit, P., Pécontal, E., & Binette, L. 2002, ApJ, 568, 627
- Crenshaw, D. M., & Kraemer, S. B. 2000*a*, ApJ, 532, L101
- \_\_\_\_\_ 2000*b*, ApJ, 532, 247
- \_\_\_\_\_ 2005, ApJ, 625, 680
- Crenshaw, D. M., Kraemer, S. B., & George, I. M. 2003, ARA&A, 41, 117
- Das, V., Crenshaw, D. M., Kraemer, S. B., & Deo, R. P. 2006, AJ, 132, 620
- Dopita, M. A., & Sutherland, R. S. 1996, ApJS, 102, 161
- Dopita, M. A., et al. 1997, ApJ, 490, 202
- Emsellem, E., Fathi, K., Wozniak, H., Ferruit, P., Mundell, C. G., & Schinnerer, E. 2006, MNRAS, 365, 367
- Evans, I. N., Ford, H. C., Kinney, A. L., Antonucci, R. R. J., Armus, L., & Caganoff, S. 1991, ApJ, 369, L27
- Ferguson, J. W., Ferland, G. J., & Pradhan, A. K. 1995, ApJ, 438, L55
- Gerssen, J., Allington-Smith, J., Miller, B. W., Turner, J. E. H., & Walker, A. 2006, MNRAS, 365, 29
- Grimes, J. P., Kriss, G. A., & Espey, B. R. 1999, ApJ, 526, 130
- Groves, B. A., Cecil, G., Ferruit, P., & Dopita, M. A. 2004, ApJ, 611, 786
- Inglis, M. D. Young, S., Hough, J. H., Gledhill, T., Axon, D. J., Bailey, J. A., & Ward, M. J. 1995, MNRAS, 275, 398
- Kinkhabwala, A, et al. 2002, ApJ, 575, 732
- Kraemer, S. B., & Crenshaw, D. M. 2000*a*, ApJ, 532, 256
- \_\_\_\_\_ 2000*b*, ApJ, 544, 763
- Kriss, G. A. 1994, in *Astronomical Data Analysis Software and Systems III*, eds. D. R. Crabtree, R. J. Hanisch, & J. Barnes, (*A. S. P. Conf. Series 61*, ASP, San Francisco), 437

- Kriss, G. A., Davidsen, A. F., Blair, W. P., Ferguson, H. C., & Long, K. S. 1992, *ApJ*, 394, L37
- Kriss, G. A., Blustin, A., Branduardi-Raymont, G., Green, R. F., Hutchings, J., & Kaiser, M. E. 2003, *Å*, 403, 473
- Moos, H. W., et al. 2000, *ApJ*, 538, L1
- Morganti, R., Oosterloo, T., & Tsvetanov, Z. 1998, *AJ*, 115, 915
- Morse, J. A., Raymond, J. C., & Wilson, A. S. 1996, *PASP*, 108, 426
- Sutherland, R. S., Bicknell, G. V., & Dopita, M. A. 1993, *ApJ*, 414, 510
- Unger, S. W., Lewis, J. R., Pedlar, A., & Axon, D. J. 1992, *MNRAS*, 258, 371
- Wilson, A. S., & Ulvestad, J. S. 1983, *ApJ*, 275, 8
- Young, A. J., Wilson, A. S., Shopbell, P. L. 2001, *ApJ*, 556, 6



Table 1. FUSE Observations of NGC 1068

| Data Set<br>(Pointing) | Start Time<br>UT     | Orbit | Integration Time<br>(sec) | Aperture | Peak-up    |
|------------------------|----------------------|-------|---------------------------|----------|------------|
| P1110202000            | 2001-Nov-28 09:57:00 | 8     | 21951                     | LWRS     | MDRS       |
| A1390201000            | 2001-Nov-29 01:23:46 | 9     | 19541                     | HIRS     | MDRS, HIRS |
| A1390202000            | 2001-Nov-29 16:34:46 | 9     | 19828                     | HIRS     | MDRS, HIRS |
| A1390203000            | 2001-Nov-30 08:38:00 | 7     | 18023                     | HIRS     | MDRS, HIRS |
| A1390204000            | 2001-Nov-30 20:53:34 | 9     | 17962                     | HIRS     | MDRS, HIRS |
| A1390205000            | 2001-Dec-01 11:52:20 | 8     | 19080                     | HIRS     | MDRS, HIRS |

Table 2. Flux Normalization Factor

| Position | LiF1A | LiF2A | SiC1B | SiC2B | STIS |
|----------|-------|-------|-------|-------|------|
| A        | 1.00  | 1.51  | 0.86  | 2.48  | 2.50 |
| B        | 1.00  | 1.42  | 0.97  | 1.70  | 2.03 |
| C        | 1.00  | 1.17  | 1.03  | 1.07  | 1.74 |
| D        | 1.00  | 1.29  | 0.96  | 1.29  | 2.28 |
| E        | 1.00  | 1.15  | 0.78  | 0.87  | 3.95 |
| F        | 1.00  | 1.10  | 1.28  | 1.17  | 4.84 |
| G        | 1.00  | 1.51  | 2.75  | 1.83  | 6.05 |

Table 3. UV Emission Lines in Spectrum with Large LWRs Aperture

| Emission Line | Å       | Flux <sup>a</sup><br>( $10^{-14}$ ergs s <sup>-1</sup> cm <sup>-2</sup> ) | FWHM<br>(km s <sup>-1</sup> ) | Velocity <sup>b</sup><br>(km s <sup>-1</sup> ) |
|---------------|---------|---------------------------------------------------------------------------|-------------------------------|------------------------------------------------|
| C III Narrow  | 977.02  | $46.2 \pm 4.0$                                                            | $720 \pm 52$                  | $28 \pm 16$                                    |
| C III Broad   |         | $10.1 \pm 5.3$                                                            | $3489 \pm 1626$               | $1047 \pm 544$                                 |
| N III Narrow  | 990.98  | $19.8 \pm 5.2$                                                            | $934 \pm 177$                 | $381 \pm 47$                                   |
| N III Broad   |         | $28.5 \pm 6.5$                                                            | $3393 \pm 315$                | $1371 \pm 431$                                 |
| Ly $\beta$    | 1025.72 | $28.4 \pm 6.5$                                                            | $3392 \pm 314$                | $1340 \pm 46$                                  |
| O VI Narrow r | 1037.63 | $41.1 \pm 2.5$                                                            | $390 \pm 31$                  | $290 \pm 6$                                    |
| O VI Narrow b |         | $59.7 \pm 3.7$                                                            | $340 \pm 6$                   | $76 \pm 4$                                     |
| O VI Broad    | 1034.00 | $338 \pm 8.1$                                                             | $3584 \pm 184$                | $52 \pm 61$                                    |
| He II Narrow  | 1085.15 | $45 \pm 2$                                                                | $993 \pm 44$                  | $167 \pm 16$                                   |
| He II Broad   |         | $14.4 \pm 3.2$                                                            | $4286 \pm 235$                | $-3912 \pm 354$                                |

<sup>a</sup>With  $E_{B-V} = 0.00$ .

<sup>b</sup>With respect to the systemic redshift  $z = 0.0038$  km s<sup>-1</sup>.

Table 4. UV Emission Lines at Position A-G

| Line               | $\text{\AA}$ | Flux <sup>a</sup><br>( $10^{-14} \text{ ergs s}^{-1} \text{ cm}^{-2}$ ) | FWHM <sup>b</sup><br>( $\text{km s}^{-1}$ ) | Velocity <sup>b,c</sup><br>( $\text{km s}^{-1}$ ) | Comment |
|--------------------|--------------|-------------------------------------------------------------------------|---------------------------------------------|---------------------------------------------------|---------|
| Position A         |              |                                                                         |                                             |                                                   |         |
| C III              | 977.02       | $2.3 \pm 1.5$                                                           | $564 \pm 243$                               | $604 \pm 198$                                     |         |
| N III Narrow       | 990.98       | $2.0 \pm 1.0$                                                           | $230 \pm 108$                               | $140 \pm 60$                                      |         |
| N III Broad        |              | $0.0 \pm 0.0$                                                           | $25434 \pm 92630$                           | $1284 \pm 30138$                                  |         |
| Ly $\beta$         | 1025.72      | $0.7 \pm 0.4$                                                           | $878 \pm 0$                                 | $257 \pm 192$                                     |         |
| O VI Narrow        | 1037.63      | $7.9 \pm 0.6$                                                           | $655 \pm 38$                                | $376 \pm 6$                                       |         |
| O VI Broad         | 1034.00      | $7.8 \pm 0.8$                                                           | $3131 \pm 488$                              | $1192 \pm 207$                                    |         |
| He II              | 1085.15      | $3.1 \pm 1.0$                                                           | $1098 \pm 968$                              | $16 \pm 116$                                      |         |
| Ly $\alpha$ Narrow | 1215.67      | $37.1 \pm 2.8$                                                          | $1580 \pm 96$                               | $393 \pm 37$                                      |         |
| Ly $\alpha$ Broad  |              | $19.2 \pm 2.1$                                                          | $5283 \pm 743$                              | $393 \pm 0$                                       |         |
| C IV Narrow        | 1549.50      | $18.8 \pm 6.5$                                                          | $1432 \pm 95$                               | $374 \pm 96$                                      |         |
| C IV Broad         |              | $7.6 \pm 1.4$                                                           | $6178 \pm 1623$                             | $507 \pm 424$                                     |         |
| He II              | 1640.46      | $9.8 \pm 0.7$                                                           | $1858 \pm 134$                              | $10 \pm 67$                                       |         |
| N III              | 1750.00      | $0.3 \pm 0.5$                                                           | $1818 \pm 1916$                             | $3691 \pm 1758$                                   |         |
| C III] Narrow      | 1908.73      | $6.8 \pm 1.3$                                                           | $2111 \pm 228$                              | $43 \pm 72$                                       |         |
| C III] Broad       |              | $9.4 \pm 1.3$                                                           | $6883 \pm 1106$                             | $-535 \pm 348$                                    |         |
| Position B         |              |                                                                         |                                             |                                                   |         |
| C III              | 977.02       | $14.1 \pm 3.5$                                                          | $596 \pm 290$                               | $-31 \pm 55$                                      |         |
| N III Narrow       | 990.98       | $8.8 \pm 3.5$                                                           | $1048 \pm 351$                              | $519 \pm 190$                                     |         |
| N III Broad        |              | $2.4 \pm 2.7$                                                           | $2005 \pm 2083$                             | $2332 \pm 1454$                                   |         |
| Ly $\beta$         | 1025.72      | $5.2 \pm 1.0$                                                           | $878 \pm 0$                                 | $-2 \pm 105$                                      |         |
| O VI Narrow        | 1037.63      | $23.9 \pm 5.1$                                                          | $385 \pm 34$                                | $333 \pm 12$                                      |         |
| O VI Broad         | 1034.00      | $123.5 \pm 2.5$                                                         | $2535 \pm 35$                               | $298 \pm 49$                                      |         |
| He II              | 1085.15      | $10.2 \pm 1.5$                                                          | $1108 \pm 177$                              | $-18 \pm 74$                                      |         |
| Ly $\alpha$ Narrow | 1215.67      | $158.2 \pm 0.4$                                                         | $1528 \pm 18$                               | $366 \pm 10$                                      |         |
| Ly $\alpha$ Broad  |              | $119.9 \pm 7.8$                                                         | $7676 \pm 787$                              | $366 \pm 0$                                       |         |
| C IV Narrow        | 1549.50      | $95.8 \pm 2.9$                                                          | $1348 \pm 32$                               | $284 \pm 12$                                      |         |
| C IV Broad         |              | $60.3 \pm 3.1$                                                          | $4424 \pm 172$                              | $-92 \pm 66$                                      |         |
| He II              | 1640.46      | $47.0 \pm 1.6$                                                          | $1872 \pm 66$                               | $-234 \pm 31$                                     |         |
| N III              | 1750.00      | $4.1 \pm 1.0$                                                           | $1848 \pm 569$                              | $299 \pm 169$                                     |         |
| C III] Narrow      | 1908.73      | $29.6 \pm 2.4$                                                          | $2229 \pm 95$                               | $-126 \pm 27$                                     |         |
| C III] Broad       |              | $36.7 \pm 2.2$                                                          | $6296 \pm 408$                              | $-678 \pm 110$                                    |         |
| Position C         |              |                                                                         |                                             |                                                   |         |
| C III              | 977.02       | $38.6 \pm 4.3$                                                          | $911 \pm 99$                                | $-7 \pm 49$                                       |         |
| N III Narrow       | 990.98       | $11.0 \pm 10.3$                                                         | $989 \pm 558$                               | $-8 \pm 202$                                      |         |
| N III Broad        |              | $29.0 \pm 14.8$                                                         | $3207 \pm 767$                              | $870 \pm 770$                                     |         |

Table 4—Continued

| Line               | Å       | Flux <sup>a</sup><br>( $10^{-14}$ ergs s <sup>-1</sup> cm <sup>-2</sup> ) | FWHM <sup>b</sup><br>(km s <sup>-1</sup> ) | Velocity <sup>b,c</sup><br>(km s <sup>-1</sup> ) | Comment |
|--------------------|---------|---------------------------------------------------------------------------|--------------------------------------------|--------------------------------------------------|---------|
| Ly $\beta$         | 1025.72 | $12.6 \pm 2.9$                                                            | $878 \pm 132$                              | $-84 \pm 61$                                     |         |
| O VI Narrow        | 1037.63 | $46.2 \pm 6.3$                                                            | $333 \pm 22$                               | $290 \pm 9$                                      |         |
| O VI Broad         | 1034.00 | $325.3 \pm 43.4$                                                          | $2523 \pm 118$                             | $162 \pm 55$                                     |         |
| He II              | 1085.15 | $17.0 \pm 1.7$                                                            | $1149 \pm 116$                             | $-70 \pm 55$                                     |         |
| Ly $\alpha$ Narrow | 1215.67 | $278.5 \pm 4.7$                                                           | $1352 \pm 21$                              | $395 \pm 7$                                      |         |
| Ly $\alpha$ Broad  |         | $253.3 \pm 13.5$                                                          | $9104 \pm 641$                             | $395 \pm 0$                                      |         |
| C IV Narrow        | 1549.50 | $167.6 \pm 9.9$                                                           | $1472 \pm 59$                              | $172 \pm 19$                                     |         |
| C IV Broad         |         | $131.0 \pm 9.8$                                                           | $4193 \pm 244$                             | $-390 \pm 73$                                    |         |
| He II              | 1640.46 | $92.0 \pm 3.8$                                                            | $1830 \pm 82$                              | $-406 \pm 35$                                    |         |
| N III              | 1750.00 | $10.1 \pm 1.6$                                                            | $2605 \pm 456$                             | $118 \pm 160$                                    |         |
| C III] Narrow      | 1908.73 | $38.4 \pm 4.4$                                                            | $1902 \pm 107$                             | $-270 \pm 28$                                    |         |
| C III] Broad       |         | $80.3 \pm 4.0$                                                            | $5053 \pm 219$                             | $-672 \pm 56$                                    |         |
| Position D         |         |                                                                           |                                            |                                                  |         |
| C III              | 977.02  | $31.2 \pm 3.2$                                                            | $900 \pm 0$                                | $183 \pm 52$                                     |         |
| N III Narrow       | 990.98  | $11.1 \pm 3.3$                                                            | $888 \pm 243$                              | $372 \pm 93$                                     |         |
| N III Broad        |         | $44.1 \pm 71.1$                                                           | $5775 \pm 3455$                            | $-3804 \pm 4077$                                 |         |
| Ly $\beta$         | 1025.72 | $6.2 \pm 1.8$                                                             | $878 \pm 0$                                | $-46 \pm 0$                                      |         |
| O VI Narrow        | 1037.63 | $14.9 \pm 0.9$                                                            | $380 \pm 31$                               | $287 \pm 9$                                      |         |
| O VI Broad         | 1034.00 | $181.8 \pm 17.1$                                                          | $2506 \pm 126$                             | $-283 \pm 32$                                    |         |
| He II              | 1085.15 | $15.6 \pm 1.2$                                                            | $940 \pm 85$                               | $21 \pm 36$                                      |         |
| Ly $\alpha$ Narrow | 1215.67 | $291.4 \pm 14.3$                                                          | $1248 \pm 42$                              | $388 \pm 7$                                      |         |
| Ly $\alpha$ Broad  |         | $145.9 \pm 10.6$                                                          | $5718 \pm 1708$                            | $388 \pm 0$                                      |         |
| C IV Narrow        | 1549.50 | $143.7 \pm 18.7$                                                          | $1777 \pm 125$                             | $6 \pm 21$                                       |         |
| C IV Broad         |         | $105.9 \pm 18.4$                                                          | $3388 \pm 130$                             | $-793 \pm 17$                                    |         |
| He II              | 1640.46 | $88.6 \pm 3.4$                                                            | $1928 \pm 73$                              | $-504 \pm 35$                                    |         |
| N III              | 1750.00 | $9.1 \pm 2.3$                                                             | $2415 \pm 404$                             | $15 \pm 188$                                     |         |
| C III] Narrow      | 1908.73 | $45.3 \pm 2.9$                                                            | $2181 \pm 493$                             | $-367 \pm 114$                                   |         |
| C III] Broad       |         | $58.8 \pm 0.0$                                                            | $5201 \pm 0$                               | $-877 \pm 39538$                                 |         |
| Position E         |         |                                                                           |                                            |                                                  |         |
| C III              | 977.02  | $17.7 \pm 4.2$                                                            | $1149 \pm 412$                             | $27 \pm 119$                                     |         |
| N III Narrow       | 990.98  | $6.7 \pm 2.6$                                                             | $577 \pm 177$                              | $375 \pm 84$                                     |         |
| N III Broad        |         | $12.9 \pm 15.0$                                                           | $2584 \pm 2236$                            | $-764 \pm 1615$                                  |         |
| Ly $\beta$         | 1025.72 | $3.6 \pm 1.2$                                                             | $878 \pm 0$                                | $68 \pm 0$                                       |         |
| O VI Narrow        | 1037.63 | $13.0 \pm 1.2$                                                            | $639 \pm 42$                               | $252 \pm 17$                                     |         |
| O VI Broad         | 1034.00 | $52.9 \pm 2.9$                                                            | $3145 \pm 168$                             | $-475 \pm 78$                                    |         |
| He II              | 1085.15 | $12.1 \pm 0.8$                                                            | $818 \pm 58$                               | $208 \pm 25$                                     |         |
| Ly $\alpha$ Narrow | 1215.67 | $359.0 \pm 9.4$                                                           | $1265 \pm 24$                              | $474 \pm 7$                                      |         |

Table 4—Continued

| Line               | $\text{\AA}$ | Flux <sup>a</sup><br>( $10^{-14} \text{ ergs s}^{-1} \text{ cm}^{-2}$ ) | FWHM <sup>b</sup><br>( $\text{km s}^{-1}$ ) | Velocity <sup>b,c</sup><br>( $\text{km s}^{-1}$ ) | Comment |
|--------------------|--------------|-------------------------------------------------------------------------|---------------------------------------------|---------------------------------------------------|---------|
| Ly $\alpha$ Broad  |              | $72.4 \pm 6.4$                                                          | $4953 \pm 1166$                             | $474 \pm 0$                                       |         |
| C IV Narrow        | 1549.50      | $107.2 \pm 10.9$                                                        | $1812 \pm 141$                              | $79 \pm 71$                                       |         |
| C IV Broad         |              | $63.5 \pm 11.0$                                                         | $3241 \pm 134$                              | $-981 \pm 99$                                     |         |
| He II              | 1640.46      | $64.0 \pm 2.9$                                                          | $4490 \pm 157$                              | $1112 \pm 5$                                      |         |
| N III              | 1750.00      | $9.1 \pm 1.6$                                                           | $2986 \pm 549$                              | $362 \pm 237$                                     |         |
| C III] Narrow      | 1908.73      | $30.3 \pm 4.2$                                                          | $2185 \pm 135$                              | $82 \pm 55$                                       |         |
| C III] Broad       |              | $39.3 \pm 4.3$                                                          | $4489 \pm 251$                              | $-1026 \pm 162$                                   |         |
| Position F         |              |                                                                         |                                             |                                                   |         |
| C III              | 977.02       | $15.8 \pm 4.3$                                                          | $900 \pm 270$                               | $424 \pm 119$                                     |         |
| N III Narrow       | 990.98       | $7.1 \pm 5.8$                                                           | $1056 \pm 625$                              | $164 \pm 289$                                     |         |
| N III Broad        |              | $8.0 \pm 9.4$                                                           | $3540 \pm 1540$                             | $333 \pm 1681$                                    |         |
| Ly $\beta$         | 1025.72      | $9.0 \pm 0.7$                                                           | $878 \pm 0$                                 | $257 \pm 47$                                      |         |
| O VI Narrow        | 1037.63      | $11.4 \pm 1.9$                                                          | $219 \pm 30$                                | $152 \pm 17$                                      |         |
| O VI Broad         | 1034.00      | $96.1 \pm 15.6$                                                         | $1785 \pm 16$                               | $456 \pm 20$                                      |         |
| He II              | 1085.15      | $8.3 \pm 0.7$                                                           | $665 \pm 66$                                | $266 \pm 25$                                      |         |
| Ly $\alpha$ Narrow | 1215.67      | $299.6 \pm 9.6$                                                         | $1223 \pm 33$                               | $520 \pm 10$                                      |         |
| Ly $\alpha$ Broad  |              | $46.8 \pm 7.4$                                                          | $4582 \pm 1014$                             | $520 \pm 0$                                       |         |
| C IV Narrow        | 1549.50      | $73.3 \pm 5.9$                                                          | $1181 \pm 72$                               | $313 \pm 29$                                      |         |
| C IV Broad         |              | $34.7 \pm 5.3$                                                          | $3365 \pm 323$                              | $-461 \pm 178$                                    |         |
| He II              | 1640.46      | $42.6 \pm 2.8$                                                          | $1132 \pm 74$                               | $305 \pm 33$                                      |         |
| N III              | 1750.00      | $4.7 \pm 1.6$                                                           | $1289 \pm 702$                              | $1040 \pm 211$                                    |         |
| C III] Narrow      | 1908.73      | $29.1 \pm 1.2$                                                          | $1500 \pm 77$                               | $400 \pm 41$                                      |         |
| C III] Broad       |              | $14.3 \pm 0.9$                                                          | $3848 \pm 392$                              | $-1679 \pm 139$                                   |         |
| Position G         |              |                                                                         |                                             |                                                   |         |
| C III              | 977.02       | $12.5 \pm 5.4$                                                          | $344 \pm 133$                               | $91 \pm 80$                                       |         |
| N III Narrow       | 990.98       | $8.7 \pm 4.2$                                                           | $833 \pm 398$                               | $417 \pm 148$                                     |         |
| N III Broad        |              | $10.8 \pm 58.4$                                                         | $3237 \pm 7653$                             | $-2085 \pm 9897$                                  |         |
| Ly $\beta$         | 1025.72      | $4.3 \pm 0.9$                                                           | $878 \pm 0$                                 | $-61 \pm 93$                                      |         |
| O VI Narrow        | 1037.63      | $13.3 \pm 2.6$                                                          | $372 \pm 16$                                | $189 \pm 14$                                      |         |
| O VI Broad         | 1034.00      | $37.6 \pm 4.9$                                                          | $2852 \pm 422$                              | $416 \pm 95$                                      |         |
| He II              | 1085.15      | $7.5 \pm 1.4$                                                           | $617 \pm 105$                               | $150 \pm 58$                                      |         |
| Ly $\alpha$ Narrow | 1215.67      | $152.2 \pm 5.8$                                                         | $1208 \pm 35$                               | $523 \pm 12$                                      |         |
| Ly $\alpha$ Broad  |              | $32.2 \pm 3.9$                                                          | $4627 \pm 848$                              | $523 \pm 0$                                       |         |
| C IV Narrow        | 1549.50      | $50.8 \pm 3.0$                                                          | $1227 \pm 62$                               | $251 \pm 27$                                      |         |
| C IV Broad         |              | $19.5 \pm 2.3$                                                          | $4877 \pm 581$                              | $-283 \pm 216$                                    |         |
| He II              | 1640.46      | $28.0 \pm 1.7$                                                          | $1047 \pm 58$                               | $265 \pm 29$                                      |         |
| N III              | 1750.00      | $1.5 \pm 0.8$                                                           | $708 \pm 334$                               | $1453 \pm 195$                                    |         |

Table 4—Continued

| Line          | $\text{\AA}$ | Flux <sup>a</sup><br>( $10^{-14}$ ergs s <sup>-1</sup> cm <sup>-2</sup> ) | FWHM <sup>b</sup><br>(km s <sup>-1</sup> ) | Velocity <sup>b,c</sup><br>(km s <sup>-1</sup> ) | Comment |
|---------------|--------------|---------------------------------------------------------------------------|--------------------------------------------|--------------------------------------------------|---------|
| C III] Narrow | 1908.73      | $12.5 \pm 3.1$                                                            | $1797 \pm 252$                             | $333 \pm 83$                                     |         |
| C III] Broad  |              | $12.4 \pm 3.5$                                                            | $4491 \pm 658$                             | $-870 \pm 510$                                   |         |

<sup>a</sup>With  $E_{B-V} = 0.00$ .

<sup>b</sup>Values with zero errors are pre-fixed.

<sup>c</sup>With respect to the systemic redshift  $z = 0.0038$  km s<sup>-1</sup>.

### Figure Captions

Fig. 1.— HST image of the central region of NGC 1068 obtained with WFPC2/F218W, with a field of view of  $\sim 25''$ . The north is up, east is to the left, and the nucleus is marked with a cross. Seven FUSE slit positions are marked with letters A-G, while C is at the nucleus. Note that the actual slit width is slightly wider than that in the illustration. The narrow slits along the near-perpendicular direction are that of HST/STIS: The three parallel slits along the northeast and southwest direction are  $0''.2$  wide (STIS-A, Cecil et al. 2002), and the single long slit is another that is  $0''.1$  wide (STIS-B, Crenshaw & Kraemer 2000b).

Fig. 2.— FUSE spectrum of NGC 1068 at full aperture ( $30''$  square). The data are binned to  $0.1 \text{ \AA}$ . Propagation errors are plotted in the lower panel. The effect of major geocoronal emission lines are marked with the Earth symbols, and data at these wavelength bins are removed.

Fig. 3.— Typical relative drifts of the FUSE SiC2 channel vs. orbital time (after orbital sunset), with respect to the LiF1 channel. Data are collected from various orbits with occultation.

Fig. 4.— Count rate vs. orbital time in a sample data set A1390201001. Different channels register their peaks at different times, which are presumed to coincide with the nucleus position. Incomplete segments are not used (Position A in SiC channels and position G in LiF channels).

Fig. 5.— Merged FUSE spectra at seven FUSE slit positions, binned to  $0.5 \text{ \AA}$ . The nucleus is at position C.

Fig. 6.— STIS spectra extracted at different windows that correspond to the respective FUSE slits. Fluxes are normalized to the corresponding FUSE data.

Fig. 7.— Line profiles at different FUSE slit positions. The lines are redshifted O VI  $\lambda 1037$  (centered at  $0 \text{ km s}^{-1}$ ), C IV  $\lambda 1549$ , N III  $\lambda 991$ , and C III  $\lambda 977$ , respectively.

Fig. 8.— Intensities of major emission lines at different FUSE positions, from STIS (left panels) and FUSE (right panels). When a line is fitted with multiple components with reasonably accuracy, the flux presented is their sum. For C III  $\lambda 977$ , only one narrow component is fitted.

Fig. 9.— Emission and absorption lines near the O VI wavelength, from position A to G after subtraction of the fitted continuum. The spectrum marked “All” in the lower bottom



panel is from the data with a large LWRs aperture. The dashed lines mark the wavelengths of redshifted Ly $\beta$ , O VI  $\lambda\lambda$ 1031.95/1037.63, respectively, from left to right.

Fig. 10.— Properties of fitted O VI emission line at different FUSE slit positions. Line fluxes are in units of  $10^{-14}$  ergs s $^{-1}$  cm $^{-2}$ , FWHM and velocity are in units of km s $^{-1}$ .

Fig. 11.— Properties of fitted C IV emission line (from HST/STIS) at different FUSE positions. Line fluxes are in units of  $10^{-14}$  ergs s $^{-1}$  cm $^{-2}$ , FWHM and velocity are in units of km s $^{-1}$ .

Fig. 12.— Line ratios at different slit positions. The fitting results of C IV and O VI are for their narrow-line components.




Fig. 1.—




Fig. 2.—




Fig. 3.—




Fig. 4.—




Fig. 5.—




Fig. 6.—




Fig. 7.—






Fig. 8.—




Fig. 9.—




Fig. 10.—




Fig. 11.—




Fig. 12.—